Table 1-1: Global atmospheric concentration (ppm unless otherwise specified), rate of concentration change (ppb/year) and

atmospheric lifetime (years) of selected greenhouse gases

Atmospheric Variable	CO <sub>2</sub>	CH <sub>4</sub>	N <sub>2</sub> O	SF <sub>6</sub> <sup>a</sup>	CF <sub>4</sub> <sup>a</sup>
Pre-industrial atmospheric	280	0.722	0.270	0	40
concentration					
Atmospheric concentration <sup>b</sup>	370.3	1.842	0.316	4.7	80
Rate of concentration change <sup>c</sup>	1.5 <sup>d</sup>	$0.007^{d}$	0.0008	0.24	1.0
Atmospheric Lifetime	$5-200^{\rm e}$	12 <sup>f</sup>	114 <sup>f</sup>	3,200	>50,000

Source: Current atmospheric concentrations for CO<sub>2</sub>, CH<sub>4</sub>, N<sub>2</sub>O, and SF<sub>6</sub> are from Blasing and Jones (2002). All other data is from IPCC

<sup>&</sup>lt;sup>a</sup> Concentrations in parts per trillion (ppt) and rate of concentration change in ppt/year.

<sup>&</sup>lt;sup>b</sup> Concentration for CO<sub>2</sub> was measured in 2001. Concentrations for all other gases were measured in 2000.

<sup>&</sup>lt;sup>c</sup> Rate is calculated over the period 1990 to 1999.

<sup>&</sup>lt;sup>d</sup> Rate has fluctuated between 0.9 and 2.8 ppm per year for CO<sub>2</sub> and between 0 and 0.013 ppm per year for CH<sub>4</sub> over the period 1990 to 1999. 
<sup>e</sup> No single lifetime can be defined for CO<sub>2</sub> because of the different rates of uptake by different removal processes. 
<sup>f</sup> This lifetime has been defined as an "adjustment time" that takes into account the indirect effect of the gas on its own residence time.